

Koivula M., Vanha-Majamaa I. (2021). Eri hakkuu- ja luonnonhoitomenetelmien vaikutukset monimuotoisuuteen Fennoskandiassa. *Metsätieteen aikakauskirja* 2021-10481. <https://doi.org/10.14214/ma.10481>.

## **Liitetiedosto L1**

### **Artikkelissa käytetyt kokeellisen tutkimuksen kirjallisuuslähteet**

Järjestys: (1) koeasetelman nimi, (2) ensimmäinen kirjoittaja.

#### **Lahopuun lisäyskoe**

Pasanen, H. 2017: Ecological effects of disturbance-based restoration in boreal forests. *Väitöskirja*, Itä-Suomen yliopisto.

Pasanen, H., Junninen, K. & Kouki, J. 2014: Restoring dead wood in forests diversifies wood-decaying fungal assemblages but does not quickly benefit red-listed species. *For. Ecol. Manag.* 312: 92-100.

#### **DISTDYN**

Jokela, J., Juutilainen, K., Korpela, L., Kouki, J., Kuntsi, S., Koivula, M. & Siitonen, J. 2018: Cross-taxon congruence and relationships to stand characteristics of vascular plants, bryophytes, polyporous fungi and beetles in mature managed boreal forests. *Ecol. Ind.* 85: 137-145.

Jokela, J., Siitonen, J. & Koivula, M. 2019: Short-term effects of selection, gap, patch and clear cutting on the beetle fauna in boreal spruce-dominated forests. *For Ecol Manag* 446: 29-37.

Koivula, M., Kuuluvainen, T., Hallman, E., Kouki, J., Siitonen, J. & Valkonen, S. 2014: Forest management inspired by natural disturbance dynamics (DISTDYN) – a long-term research and development project in Finland. *Scand. J. For. Res.* 29: 579-592.

Koivula, M., Hallman, E., Kouki, J., Kuuluvainen, T., Siitonen, J. & Valkonen, S. 2012: Luonnonmetsän inspiroimaa metsänhoitoa tutkitaan aluetason koejärjestelyllä. *Metsätieteen Aikakauskirja* 1/2012: 23-31.

Pasanen, H., Juutilainen, K. & Siitonen, J. 2019: Responses of polypore fungi following disturbance-emulating harvesting treatments and deadwood creation in boreal Norway spruce dominated forests. *Scand. J. For. Res.* 34: 557-568.

#### **Effaråsen**

Djupström, L. & Weslien, J. 2019: Effaråsen – att bruka och bevara i gammal tallskog. *SkogForsk* Arbetsrapport 1009-2019.

Santaniello, F. 2017: Impact of tree retention on wood production, biodiversity conservation and carbon stock changes in boreal pine forest. Väitöskirja, Sveriges Lantbruksuniversitet, Uppsala.

Santaniello, F., Djupström, L.B., Ranius, T., Rudolphi, J., Widenfalk, O. & Weslien, J. 2016: Effects of partial cutting on logging productivity, economic returns and dead wood in boreal pine forest. *For. Ecol. Manag.* 365: 152-158.

Santaniello, F., Djupström, L.B., Ranius, T., Weslien, J., Rudolphi, J. & Thor, G. 2017a: Large proportion of wood dependent lichens in boreal pine forest are confined to old hard wood. *Biodiv. Conserv.* 26: 1295-1310.

Santaniello, F., Djupström, L.B., Ranius, T., Weslien, J., Rudolphi, J. & Sonesson, J. 2017b: Simulated long-term effects of varying tree retention on wood production, dead wood and carbon stock changes. *J. Environ. Manag.* 201: 37-44.

### **Elimyssalo**

Hekkala, A.-M., Päätalo, M.-L., Tarvainen, O. & Tolvanen, A. 2014a: Restoration of young forests in eastern Finland: benefits for saproxylic beetles (Coleoptera). *Restoration Ecology* 22: 151-159.

Hekkala, A.-M., Tarvainen, O. & Tolvanen, A. 2014b: Dynamics of understory vegetation after restoration of natural characteristics in the boreal forests in Finland. *For. Ecol. Manag.* 330: 55-66.

Hekkala, A.-M., Ahtikoski, A., Päätalo, M.-L., Tarvainen, O., Siipilehto, J. & Tolvanen, A. 2016: Restoring volume, diversity and continuity of deadwood in boreal forests. *Biodivers. Conserv.* 25: 1107-1132.

Hägglund, R., Hekkala, A.-M., Hjältén, J. & Tolvanen, A. 2015: Positive effects of ecological restoration on rare and threatened flat bugs (Heteroptera: Aradidae). *J. Insect Conserv.* 19: 1089-1099.

### **Eriksköp**

Drössler, L. 2016: Heterogen skog – för komplex för skogsbruk och tillämpad skogsforskning? Rön för Sveriges Lantbruksuniversitet Nr 9.

### **EVO**

Berglund, H., Jönsson, M.T., Penttilä, R. & Vanha-Majamaa, I. 2011: The effects of burning and dead-wood creation on the diversity of pioneer wood-inhabiting fungi in managed boreal spruce forests. *For. Ecol. Manag.* 261: 1293-1305.

Komonen, A., Kuntzi, S., Toivanen, T. & Kotiaho, J.S. 2014: Fast but ephemeral effects of ecological restoration on forest beetle community. *Biodiv. Conserv.* 23: 1485-1507.

Lilja, S., De Chantal, M., Kuuluvainen, T., Vanha-Majamaa, I. & Puttonen, P. 2005: Restoring natural characteristics in managed Norway spruce [Picea abies (L.) Karst.] stands with partial cutting, dead wood creation and fire: immediate treatment effects. *Scand. J. For. Res.* 20: 68-78.

Rudolphi, J., Caruso, A., von Cräutlein, M., Laaka-Lindberg, S., Ryömä, R. & Berglund, H. 2011: Relative importance of thinned and clear-cut stands for bryophyte diversity on stumps. *For. Ecol. Manag.* 261: 1911-1918.

Toivanen, T. 2007: Short-term effects of forest restoration on beetle diversity. *Väitöskirja*, Jyväskylän yliopisto.

Toivanen, T. & Kotiaho, J. 2007a: Mimicking natural disturbances of boreal forests: the effects of controlled burning and creating dead wood on beetle diversity. *Biod. Conserv.* 16: 3193-3211.

Toivanen, T. & Kotiaho, J.S. 2007b: Burning of logged sites to protect beetles in managed boreal forests. *Cons. Biol.* 21: 1562-1572.

Toivanen, T. & Kotiaho, J. 2010: The preferences of saproxylic beetle species for different dead wood types created in forest restoration treatments. *Can. J. For. Res.* 40: 445-464.

Toivanen, T., Heikkilä, T. & Koivula, M.J. 2014: Emulating natural disturbances in boreal Norway spruce forests: effects on ground beetles (Coleoptera, Carabidae). *For. Ecol. Manag.* 314: 64-74.

Toivanen, T., Liikanen, V. & Kotiaho, J.S. 2009: Effects of forest restoration treatments on the abundance of bark beetles in Norway spruce forests of southern Finland. *For. Ecol. Manag.* 257: 117-125.

Toivanen, T., Selonen, V. & Kotiaho, J.S. 2004: Effects of management and restoration on forest biodiversity: an experimental approach. *EFI Proceedings No.* 51: 493-503.

Vanha-Majamaa, I., Lilja, S., Ryömä, R., Kotiaho, J.S., Laaka-Lindberg, S., Lindberg, H., Puttonen, P., Tamminen, P., Toivanen, T. & Kuuluvainen, T. 2007: Rehabilitating boreal forest structure and species composition in Finland through logging, dead wood creation and fire: the EVO experiment. *For. Ecol. Manag.* 250: 77-88.

## Fagerön

Lindhe, A. & Lindelöw, A. 2004: Cut high stumps of spruce, birch, aspen and oak as breeding substrates for saproxylic beetles. *For. Ecol. Manag.* 203: 1-20.

Lindhe, A., Åsenblad, N. & Toresson, H.-G. 2004: Cut logs and high stumps of spruce, birch, aspen and oak – nine years of saproxylic fungi succession. *Biol. Conserv.* 119: 443-454.

## FIRE

den Herder, M., Kouki, J. & Ruusila, V. 2009: The effects of timber harvest, forest fire, and herbivores on regeneration of deciduous trees in boreal pine-dominated forests. *Can. J. For. Res.* 39: 712-722.

Eales, J., Haddaway, N.R., Bernes, C., Cooke, S.J., Jonsson, B.G., Kouki, J., Petrokofsky, G. & Taylor, J.J. 2018: What is the effect of prescribed burning in temperate and boreal forest on biodiversity, beyond pyrophilous and saproxylic species? A systematic review. *Environ. Evid.* 7: UNSP 19.

Granath, G., Kouki, J., Johnson, S., Heikkala, O., Rodríguez, A. & Strengbom, J. 2018: Trade-offs in berry production and biodiversity under prescribed burning and retention regimes in boreal forests. *J. Appl. Ecol.* 55: 1658-1667.

Hämäläinen, A. 2016: Retention forestry and intensified biomass harvest: epiphytic lichen assemblages under opposing ecological effects in pine-dominated boreal forests. Väitöskirja, Itä-Suomen yliopisto.

Hämäläinen, A., Hujo, M., Heikkala, O., Junninen, K. & Kouki, J. 2016: Retention tree characteristics have major influence on the post-harvest tree mortality and availability of coarse woody debris in clear-cut areas. *For. Ecol. Manag.* 369: 66-73.

Hämäläinen, A., Kouki, J. & Löhmus, P. 2014: The value of retained Scots pines and their dead wood legacies for lichen diversity in clear-cut forests: The effects of retention level and prescribed burning. *For. Ecol. Manag.* 324: 89-100.

Heikkala, O. 2016: Emulation of natural disturbances and the maintenance of biodiversity in managed boreal forests: the effects of prescribed fire and retention forestry on insect assemblages. Väitöskirja, Itä-Suomen yliopisto.

Heikkala, O., Martikainen, P. & Kouki, J. 2017: Prescribed burning is an effective and quick method to conserve rare pyrophilous forest-dwelling flat bugs. *Insect Conserv. Div.* 10: 32-41.

Heikkala, O., Martikainen, P. & Kouki, J. 2016a: Decadal effects of emulating natural disturbances in forest management on saproxylic beetle assemblages. *Biol. Conserv.* 194: 39-47.

Heikkala, O., Seibold, S., Koivula, M., Martikainen, P., Thorn, S., Müller, J. & Kouki, J. 2016b: Retention forestry and prescribed burning result in functionally different saproxylic beetle assemblages than clear-cutting. *For. Ecol. Manag.* 359: 51-58.

Heikkala, O., Suominen, M., Junninen, K., Hämäläinen, A. & Kouki, J. 2014: Effects of retention level and fire on retention tree dynamics in boreal forests. *For. Ecol. Manag.* 328: 193-201.

Hyvärinen, E. 2006: Green-tree retention and controlled burning in restoration and conservation of beetle diversity in boreal forests. Väitöskirja, Joensuun yliopisto.

Hyvärinen, E., Kouki, J. & Martikainen, P. 2006: Fire and green-tree retention in conservation of red-listed and rare deadwood-dependent beetles in Finnish boreal forests. *Conserv. Biol.* 20: 1711-1719.

Hyvärinen, E., Kouki, J. & Martikainen, P. 2009: Prescribed fires and retention trees help to conserve beetle diversity in managed boreal forests despite their transient negative effects on some beetle groups. *Insect Conserv. Div.* 2: 93-105.

Hyvärinen, E., Kouki, J., Martikainen, P. & Lappalainen, H. 2005: Short-term effects of controlled burning and green-tree retention on beetle (Coleoptera) assemblages in managed boreal forests. *For. Ecol. Manag.* 212: 315-332.

Johnson, S., Strengbom, J. & Kouki, J. 2014: Low levels of tree retention do not mitigate the effects of clearcutting on ground vegetation dynamics. *For. Ecol. Manag.* 330: 67-74.

Junninen, K., Kouki, J. & Renvall, P. 2008: Restoration of natural legacies of fire in European boreal forests: an experimental approach to the effects on wood-decaying fungi. *Can. J. For. Res.* 38: 202-215.

Kouki, J. 2019: Fire and retention trees in facilitating biodiversity in boreal forests.  
[forest.uef.fi/jarikouki/project\\_fire.htm](http://forest.uef.fi/jarikouki/project_fire.htm)

Kouki, J. 2013: Nuoret luonnonmetsät metsien hoidon ja suojeleun mallina. Uusia mahdollisuuksia metsälännen suojeeluun talousmetsissä. Luonnon Tutkija 1-2 (2013): 4-19.

Kouki, J., Hyvärinen, E., Lappalainen, H., Martikainen, P. & Similä, M. 2012: Landscape context affects the success of habitat restoration: large-scale colonization patterns of saproxylic and fire-associated species in boreal forests. Divers. Distr. 18: 348-355.

Martikainen, P., Kouki, J. & Heikkala, O. 2006a: The effects of green tree retention and subsequent prescribed burning on ground beetles (Coleoptera: Carabidae) in boreal pine-dominated forests. Ecography 29: 659-670.

Martikainen, P., Kouki, J., Heikkala, O., Hyvärinen, E. & Lappalainen, H. 2006b: Effects of green tree retention and prescribed burning on the crown damage caused by the pine shoot beetles (*Tomicus* spp.) in pine-dominated timber harvest areas. J. Appl. Entomol. 130: 37-44.

Pitkänen, A., Kouki, J., Viiri, H. & Martikainen, P. 2008: Effects of controlled forest burning and intensity of timber harvesting on the occurrence of pine weevils, *Hylobius* spp., in regeneration areas. For. Ecol. Manag. 255: 522-529.

Pitkänen, A., Törmänen, K., Kouki, J., Järvinen, E. & Viiri, H. 2005: Effects of green tree retention, prescribed burning and soil treatment on pine weevil (*Hylobius abietis* and *Hylobius pinastri*) damage to planted Scots pine seedlings. Agr. For. Entomol. 7: 319-331.

Rodriguez Olmo, A. 2018: Promoting biodiversity and ecosystem services in managed boreal forests through disturbance-mediated functional heterogeneity. Väitöskirja, Itä-Suomen yliopisto.

Rodriguez, A. & Kouki, J. 2015: Emulating natural disturbance in forest management enhances pollination services for dominant *Vaccinium* shrubs in boreal pine-dominated forests. For. Ecol. Manag. 350: 1-12.

Rodriguez, A. & Kouki, J. 2017: Disturbance-mediated heterogeneity drives pollinator diversity in boreal managed forest ecosystems. Ecol. Appl. 27: 589-602.

Salo, K., Domisch, T. & Kouki, J. 2019: Forest wildfire and 12 years of post-disturbance succession of saprotrophic macrofungi (Basidiomycota, Ascomycota). For. Ecol. Manag. 451: 117454.

Suominen, M. 2018: Harvested and burned forests as habitats for polypore fungi. Väitöskirja, Itä-Suomen yliopisto.

Suominen, M., Junninen, K. & Kouki, J. 2019: Diversity of fungi in harvested forests 10 years after logging and burning: Polypore assemblages on different woody substrates. For. Ecol. Manag. 446: 63-70.

Suominen, M., Junninen, K., Heikkala, O. & Kouki, J. 2015: Combined effects of retention forestry and prescribed burning on polypore fungi. J. Appl. Ecol. 52: 1001-1008.

Suominen, M., Junninen, K., Heikkala, O. & Kouki, J. 2018: Burning harvested sites enhances polypore diversity on stumps and slash. For. Ecol. Manag. 414: 47-53.

Thorn, S., Seibold, S., Heikkala, O., Koivula, M., Venugopal, P. & Kouki, J. 2018: New records of Northern bats (*Eptesicus nilssonii*) in boreal clear cuts emphasize the value of green-tree retention for conservation. *Nyctalus* 19: 22-26.

### **Future Forest**

Andersson, J., Hjältén, J. & Dynesius, M. 2015: Wood-inhabiting beetles in low stumps, high stumps and logs on boreal clear-cuts: implications for dead wood management. *PLOS ONE* 10: e0118896.

Hjältén, J., Hägglund, R., Löfroth, T., Roberge, J.-M., Dynesius, M. & Olsson, J. 2017b: Forest restoration by burning and gap cutting of voluntary set-asides yield distinct immediate effects on saproxylic beetles. *Biodiv. Conserv.* 26: 1623-1640.

Hägglund, R. 2016: Ecological restoration of natural disturbances in boreal forests. *Väitöskirja*, Sveriges Lantbruksuniversitet, Umeå.

Hägglund, R. & Hjältén, J. 2018: Substrate specific restoration promotes saproxylic beetle diversity in boreal forest set-asides. *For. Ecol. Manag.* 425: 45-58.

Hägglund, R., Hekkala, A.-M., Hjältén, J. & Tolvanen, A. 2015: Positive effects of ecological restoration on rare and threatened flat bugs (Heteroptera: Aradidae). *J. Insect Conserv.* 19: 1089-1099.

Kärvemo, S., Björkman, C., Johansson, T., Weslien, J. & Hjältén, J. 2017: Forest restoration as a double-edged sword: the conflict between biodiversity conservation and pest control. *J. Appl. Ecol.* 54: 1658-1668.

### **Härjedalen**

Hagner, M. 1992: Biologiskt och ekonomiskt resultat i fältförsök med plockhuggning kombinerad med plantering. *Arbetsrapport* 63, Sveriges Lantbruksuniversitet, Umeå.

### **Hedmark**

Fossestol, K.O. & Sverdrup-Thygeson, A. 2009: Saproxylic beetles in high stumps and residual downed wood on clear-cuts and in forest edges. *Scand. J. For. Res.* 24: 403-416.

### **Leivonmäki**

Komonen, A., Halme, P., Jäntti, M., Koskela, T., Kotiaho, J. & Toivanen, T. 2014: Created substrates do not fully mimic natural substrates in restoration: the occurrence of polypores on spruce logs. *Silva Fenn.* 48: 980.

Elo, M., Halme, P., Toivanen, T. & Kotiaho, J.S. 2019: Species richness of polypores can be increased by supplementing dead wood resource into a boreal forest landscape. *J. Appl. Ecol.* 56: 1267–1277.

Norberg, A., Halme, P., Kotiaho, J.S., Toivanen, T. & Ovaskainen, O. 2019: Experimentally induced community assembly of polypores reveals the importance of both environmental filtering and assembly history. *Fungal Ecol.* 41: 137-146.

### **Medelpad-Ångermanland**

Perhans, K., Appelgren, L., Jonsson, F., Nordin, U., Söderström, B. & Gustafsson, L. 2009: Retention patches as potential refugia for bryophytes and lichens in managed forest landscapes. *Biol. Conserv.* 142: 1125-1133.

### **MONTA**

Heliölä, J., Koivula, M. & Niemelä, J. 2001: Distribution of carabid beetles (Coleoptera, Carabidae) across a boreal forest-clearcut ecotone. *Cons. Biol.* 15: 370-377.

Henttonen, H., Haukisalmi, V., Pirkkalainen, H. & Niemelä, J. 1998: Ekologisten käytävien merkityksestä keskisuomalaisissa talousmetsissä. *Metsätutkimuslaitoksen Tiedonantoja* 705: 255-271.

Jalonen, J. & Vanha-Majamaa, I. 2001: Immediate effects of four different felling methods on mature boreal spruce forest understorey vegetation in southern Finland. *For. Ecol. Manag.* 146: 25-34.

Kaila, S. (toim.) (1998) Monimuotoisuus talousmetsän uudistamisessa -hankkeen väliraportit (MONTA-hanke). *Metsätalon raportti* 62. [www.metsateho.fi/wp-content/uploads/2015/02/metsatehon\\_raportti\\_062.pdf](http://www.metsateho.fi/wp-content/uploads/2015/02/metsatehon_raportti_062.pdf)

Koivula, M. 2001: Carabid beetles (Coleoptera, Carabidae) in boreal managed forests – meso-scale ecological patterns in relation to modern forestry. *Väitöskirja*, Helsingin yliopisto.

Koivula, M. 2002a: Alternative harvesting methods and boreal carabid beetles (Coleoptera, Carabidae). *For. Ecol. Manag.* 167: 103-121.

Koivula, M. 2002b: Boreal carabid-beetle (Coleoptera, Carabidae) assemblages in thinned uneven-aged and clear-cut spruce stands. *Ann. Zool. Fenn.* 39: 131-149.

Koivula, M.J. 2012: Under which conditions does retention harvesting support ground beetles of boreal forests? *Baltic J. Coleopt.* 12: 7-26.

Koivula, M. & Niemelä, J. 2003: Gap felling as a forest harvesting method in boreal forests: responses of carabid beetles (Coleoptera, Carabidae). *Ecography* 26: 179-187.

Koivula, M.J., Venn, S., Hakola, P. & Niemelä, J. 2019: Responses of boreal ground beetles (Coleoptera, Carabidae) to different logging regimes ten years post-harvest. *For. Ecol. Manag.* 436: 27-38.

Matveinen-Huju, K. 2007: Short-term effects of variable retention on epigaeic spiders and carabid beetles in Finland. *Väitöskirja*, Helsingin yliopisto.

Matveinen-Huju, K. & Koivula, M. 2008: Effects of alternative harvesting methods on boreal forest spider assemblages. *Can. J. For. Res.* 38: 782-794.

Niemelä, J., Koivula, M. & Kotze, D.J. 2007: The effects of forestry on carabid beetles (Coleoptera: Carabidae) in boreal forests. *J. Insect Conserv.* 11: 5-18.

Pietikäinen, A.S., Haimi, J. & Siitonen, J. 2003: Short-term responses of soil macroarthropod community to clear felling and alternative forest regeneration methods. *For. Ecol. Manag.* 172: 339-353.

Pihlaja, M., Koivula, M. & Niemelä, J. 2006: Responses of boreal carabid beetle assemblages (Coleoptera, Carabidae) to clear-cutting and top-soil preparation. *For. Ecol. Manag.* 222: 182-190.

Siira-Pietikäinen, A. 2002: Decomposer community in boreal coniferous forest soil after forest harvesting. *Väitöskirja*, Jyväskylän yliopisto.

Siira-Pietikäinen, A. & Haimi, J. 2009: Changes in soil fauna 10 years after forest harvestings: Comparison between clear felling and green-tree retention methods. *For. Ecol. Manag.* 258: 332-338.

Siira-Pietikäinen, A., Pietikäinen, J.M., Fritze, H. & Haimi, J. 2001: Short-term responses of soil decomposer communities to forest management: clear felling versus alternative forest harvesting methods. *Can. J. For. Res.* 31: 88-99.

Vanha-Majamaa, I. & Jalonen, J. 2001: Green tree retention in Fennoscandian forestry. *Scand. J. For. Res. Suppl.* 3: 79-90.

Vanha-Majamaa, I., Shorohova, E., Kushnevskaya, H. & Jalonen, J. 2017: Resilience of understory vegetation after variable retention felling in boreal Norway spruce forests: A ten-year perspective. *For. Ecol. Manag.* 393: 12-28.

## NaturKultur

Drössler, L. 2016: Heterogen skog – för komplex för skogsbruk och tillämpad skogsforskning? Rön för Sveriges Lantbruksuniversitet Nr 9.

## Oak forest

Franc, N. & Götmark, F. 2008: Openness in management: Hands-off vs partial cutting in conservation forests, and the response of beetles. *Biol. Conserv.* 141: 2310-2321.

Franc, N., Götmark, F., Økland, B., Nordén, B. & Paltto, H. 2007: Factors and scales potentially important for saproxylic beetles in temperate mixed oak forest. *Biol. Conserv.* 135: 86-98.

Götmark, F., Paltto, H., Nordén, B. & Götmark, E. 2005: Evaluating partial cutting in broadleaved temperate forest under strong experimental control: short-term effects on herbaceous plants. *For. Ecol. Manag.* 214: 124-141.

Gran, O. & Götmark, F. 2019: Long-term experimental management in Swedish mixed oak-rich forests has a positive effect on saproxylic beetles after 10 years. *Biodiv. Conserv.* 28: 1451-1472.

Nordén, B., Götmark, F., Ryberg, M., Paltto, H. & Allmér, J. 2008: Partial cutting reduces species richness of fungi on woody debris in oak-rich forests. *Can. J. For. Res.* 38: 1807-1816.

- Nordén, B., Paltto, H., Claesson, C. & Götmark, F. 2012: Partial cutting can enhance epiphyte conservation in temperate oak-rich forests. *For. Ecol. Manag.* 270: 35-44.
- Økland, B., Götmark, F. & Nordén, N. 2008: Oak woodland restoration: testing the effects on biodiversity by mycetophilids in southern Sweden. *Biodiv. Cons.* 17: 2599.
- Paltto, H., Nordén, B. & Götmark, F. 2008: Partial cutting as a conservation alternative for oak (*Quercus* spp.) forest - responses of bryophytes and lichens on dead wood. *For. Ecol. Manag.* 256: 536-547.

## **PuroMONTA**

- Oldén, A., Selonen, V.A.O., Lehkonen, E. & Kotiaho, J.S. 2019a: The effect of buffer strip width and selective logging on streamside plant communities. *BMC Ecology* 19 (1): 9.
- Oldén, A., Peura, M., Saine, S., Kotiaho, J.S. & Halme, P. 2019b: The effect of buffer strip width and selective logging on riparian forest microclimate. *For. Ecol. Manag.* 453: 117623.
- Selonen, V. 2014: Forest act habitats. Finnish woodland key habitats amid the intensive forestry. Väitöskirja, Jyväskylän yliopisto.
- Selonen, V.A.O. & Kotiaho, J. 2013: Buffer strips can pre-empt extinction debt in boreal streamside habitats. *BMC Biology* 13: 24.
- Selonen, V.A.O., Mussaari, M., Toivanen, T. & Kotiaho, J.S. 2011: The conservation potential of brookside key habitats in managed boreal forests. *Silva Fenn.* 45: 1041-1052.
- Timonen, J. 2011: Woodland key habitats. A key to effective conservation of forest biodiversity? Väitöskirja, Jyväskylän yliopisto.
- Toivanen, T., Selonen, V. & Kotiaho, J.S. 2004: Effects of management and restoration on forest biodiversity: an experimental approach. *EFI Proceedings No. 51:* 493-503.

## **RETREE**

- Hautala, H. 2008: Disturbance in boreal spruce forest – immediate dynamics from stand to understorey level. Väitöskirja, Helsingin yliopisto.
- Hautala, H. & Vanha-Majamaa, I. 2006: Immediate tree uprooting after retention-felling in a coniferous boreal forest in Fennoscandia. *Can. J. For. Res.* 36: 3167-3172.
- Hautala, H., Jalonens, J., Laaka-Lindberg, S. & Vanha-Majamaa, I. 2004: Impacts of retention felling on coarse woody debris (CWD) in mature boreal spruce forests in Finland. *Biodiv. Cons.* 13: 1541-1554.
- Hautala, H., Laaka-Lindberg, S. & Vanha-Majamaa, I. 2011: Effects of retention felling on epixylic species in boreal spruce forests in southern Finland. *Rest. Ecol.* 19: 418-429.
- Matveinen-Huju, K. 2007: Short-term effects of variable retention on epigaeic spiders and carabid beetles in Finland. Väitöskirja, Helsingin yliopisto.

Matveinen-Huju, K., Koivula, M., Niemelä, J. & Rauha, A.M. 2009: Short-term effects of retention felling at mire sites on boreal spiders and carabid beetles. *For. Ecol. Manag.* 258: 2388-2398.

Matveinen-Huju, K., Niemelä, J., Rita, H. & O'Hara, R.B. 2006: Retention-tree groups in clear-cuts: do they constitute 'life-boats' for spiders and carabids? *For. Ecol. Manag.* 230: 119-135.

Vanha-Majamaa, I. & Hautala, H. 2010: Effect of the size of retention tree groups on biodiversity in forest regeneration (RETREE). Natural Resources Institute Finland (LUKE). [www.metla.fi/hanke/8532/pdf/RETREE\\_methods.pdf](http://www.metla.fi/hanke/8532/pdf/RETREE_methods.pdf)

Vanha-Majamaa, I. & Jalonen, J. 2001: Green tree retention in Fennoscandian forestry. *Scand. J. For. Res. Suppl.* 3: 79-90.

### **Rogberga**

Drössler, L. 2016: Heterogen skog – för komplex för skogsbruk och tillämpad skogsforskning? Rön för Sveriges Lantbruksuniversitet Nr 9.

### **Shelterwood**

Löhmus, P., Leppik, E., Motiejunaite, J., Suija, A. & Löhmus, A. 2012: Old selectively cut forests can host rich lichen communities - lessons from an exhaustive field survey. *Nova Hedwigia* 95: 493-515.

Löhmus, P., Löhmus, A. & Hämäläinen, A. 2018: Rapid legacy-dependent succession of lichen assemblages after forest fires: insights from two boreal regions. *J. Veg. Sci.* 29: 200-212.

### **Snöberget**

Hedenås, H. & Ericson, L. 2003: Response of epiphytic lichens on *Populus tremula* in a selective cutting experiment. *Ecol. Appl.* 13: 1124-1134.

Hedenås, H. & Ericson, L. 2008: Species occurrences at stand level cannot be understood without considering the landscape context: Cyanolichens on aspen in boreal Sweden. *Biol. Conserv.* 141: 710-718.

### **Spruce CWD**

Olsson, J., Jonsson, B.G., Hjältén, J. & Ericson, L. 2011: Addition of coarse woody debris – the early fungal succession on *Picea abies* logs in managed forests. *Biol. Conserv.* 144: 1100–1110.

Gibb, H., Pettersson, R.B., Hjältén, J., Hilszczanski, J., Ball, J.P., Johansson, T., Atlegrim, O. & Danell, K. 2006: Conservation-oriented forestry and early successional saproxylic beetles: responses of functional groups to manipulated dead wood substrates. *Biol. Conserv.* 129: 437-450.

Johansson, T., Gibb, H., Hilszczanski, J., Pettersson, R.B., Hjältén, J., Atlegrim, O., Ball, J.P. & Danell, K. 2006: Conservation-oriented manipulations of coarse woody debris affect its value as habitat for spruce-infesting bark and ambrosia beetles (Coleoptera: Scolytinae) in northern Sweden. *Can. J. For. Res.* 36: 174-185.

## UNEVEN

Hjältén, J., Joelsson, K., Gibb, H., Work, T., Löfroth, T. & Roberge, J.-M. 2017a: Biodiversity benefits for saproxyllic beetles with uneven-aged silviculture. *For. Ecol. Manag.* 402: 37-50.

Joelsson, K. 2017: Uneven-aged silviculture as a management tool to mitigate biodiversity loss. A case study of beetles in boreal forests. *Väitöskirja, Sveriges Lantbruksuniversitet, Umeå*.

Joelsson, K., Hjältén, J. & Gibb, H. 2018a: Forest management strategy affects saproxyllic beetle assemblages: a comparison of even and uneven-aged silviculture using direct and indirect sampling. *PLOS ONE* 13: e0194905.

Joelsson, K., Hjältén, J. & Work, T. 2018b: Uneven-aged silviculture can enhance within stand heterogeneity and beetle diversity. *J. Env. Manag.* 205: 1-8.

Joelsson, K., Hjältén, J., Work, T., Gibb, H., Roberge, J.-M. & Löfroth, T. 2017: Uneven-aged silviculture can reduce negative effects of forest management on beetles. *For. Ecol. Manag.* 391: 436-445.

## UPM

Rabinowitsch-Jokinen, R., Laaka-Lindberg, S. & Vanha-Majamaa, I. 2012: Immediate effects of logging, mounding and removal of logging residues on epixylic species in managed boreal Norway spruce stands in Southern Finland. *J. Sust. For.* 31: 205-229.

Rabinowitsch-Jokinen, R. & Vanha-Majamaa, I. 2010: Immediate effects of logging, mounding and removal of logging residues and stumps on coarse woody debris in managed boreal Norway spruce stands. *Silva Fenn.* 44: 51-61.